

RESEARCH ARTICLE

Effectiveness of Sage Leaf Gel on Neutrophil Count in Wound Healing

Florence Meliawaty,¹ Frita Ferlita Shafri Djohan,² Muhamad Syahrul Ramdhani³

¹Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Universitas Jenderal Achmad Yani, Cimahi, Indonesia, ²Department of Periodontology, Faculty of Dentistry, Universitas Jenderal Achmad Yani, Cimahi, Indonesia, ³Faculty of Dentistry, Universitas Jenderal Achmad Yani, Cimahi, Indonesia

Abstract

A wound involves damage to the epithelial layer of the skin, extending to the subcutis and surrounding tissues. The body initiates a healing process starting with inflammation. Neutrophils play a crucial role in this phase; excessive neutrophils can cause tissue necrosis, while insufficient neutrophils may lead to infection. Neutrophils are essential for bacterial eradication during inflammation. Sage leaf extract, known for its anti-inflammatory, antioxidant, and antibacterial properties, may aid in wound healing. This study evaluates the effectiveness of 5% sage leaf extract gel on neutrophil count during the wound healing process in Wistar rats. This study was conducted in several Universitas Jenderal Achmad Yani Cimahi laboratories from 29 May 2023 to 23 February 2024. A post-test-only control group design was used with 25 samples divided into five groups: (K-1) no treatment, tissue sampled at 13 minutes; (K-2) no treatment, tissue sampled at 103 minutes; (KN) healthy/normal control; (KP1) 5% sage leaf extract gel, tissue sampled at 13 minutes; (KP2) 5% sage leaf extract gel, tissue sampled at 103 minutes. Data analysis was performed using the Kruskal-Wallis test followed by the Mann-Whitney post hoc test ($p \leq 0.05$), indicating a statistically significant difference. The highest mean neutrophil count was observed in the KP2 group, which received 5% sage leaf extract gel and had tissue sampled at 103 minutes. The study concludes that a 5% sage leaf extract gel is effective in increasing neutrophil counts during the healing of punch wounds in Wistar rats, which contributes to accelerated wound healing.

Keywords: Neutrophil, sage, wound healing

Introduction

A wound is a disruption of tissue continuity from the epithelial layer of the skin to the subcutis layer caused by trauma.^{1,2} In dentistry, a biopsy is a standard procedure that can cause trauma to the gingival soft tissue.³ There are various types of biopsies, and in this study, a punch biopsy was used, as it is considered the most effective technique for obtaining samples for histopathological examination.⁴ When tissue is injured, it undergoes a wound-healing process that occurs in three phases, one of which is the inflammatory phase.⁵

The inflammatory phase involves the release of inflammatory mediators, including cytokines, chemokines, and phagocytic leukocytes (such as neutrophils, macrophages, platelets, and T lymphocytes). Neutrophils are a type of immune cell that moves toward the wound, which has

three main functions: the primary antimicrobial functions of phagocytosis, degranulation, and the release of nuclear material. This process works by releasing reactive oxygen species (ROS), eicosanoids, and proteolytic enzymes to kill bacteria.^{6,7} In humans, the number of neutrophils increases between the 6th and 48th hour after injury. When converted to rats, this increase occurs between the 13th and 104th minute, based on the age conversion ratio between rats and humans, which is 1:28 (1 rat day = 28 human days).^{8–10}

So far, the drug used to treat natural-based oral cavity lesions is Aloclair® gel.¹¹ Apart from Aloclair® gel, other natural ingredients can be an alternative for healing oral cavity wounds, including sage leaf extract (*Salvia officinalis* L.). Sage leaves contain flavonoids, terpenoids, and rosmarinic acid, which can act as anti-inflammatory, antibacterial, and antioxidant.^{12–14}

Copyright ©2025 by authors. This is an open access article under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (<https://creativecommons.org/licenses/by-nc-sa/4.0>).

Received: 23 October 2024; Revised: 28 April 2025; Accepted: 29 April 2025; Published: 30 April 2025

Correspondence: Florence Meliawaty, drg., M.Kes., Sp.B.M. Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Universitas Jenderal Achmad Yani. Jln. Terusan Jenderal Sudirman, Cimahi 40531, West Java, Indonesia. E-mail: florence.meliawaty@lecture.unjani.ac.id

Based on several previous studies, sage leaf extract, especially sage leaf extract with a concentration of 5%, can significantly accelerate the wound healing effect and can be considered an appropriate compound for clinical application in wound care with indicators such as an increase in the percentage of wound contraction, re-epithelialization period, and stress strength ratio.^{14,15} This study raises the question: what mechanism is enhanced that makes sage leaves considered capable of accelerating wound healing? Until now, there has been no research on the effect of sage leaf extract gel on the number of neutrophils in the wound healing process of punch rats Wistar strain.

Neutrophils were chosen to be the subject of the study because of their characteristics and functions that affect the wound healing process, such as the first body defense system that handles the wound area, the type of leukocyte with the most significant number in the human body, and its function that can phagocytize pathogenic bacteria but can also damage surrounding tissues.⁸

This study aims to evaluate the effectiveness of 5% sage leaf extract gel on the number of neutrophils during the wound healing process in Wistar strain white rats.

Methods

Laboratory experimental research design with a post-test-only control group design was used with 25 samples, which is analytic.¹³ Consists of 5 groups, namely negative group 1 (K-1): given a punch wound without being given any drug/therapy (tissue taken at the 13th minute); negative group 2 (K-2): given a punch wound without being given any drug/therapy (tissue taken at the 103rd minute); normal group (KN): without being given healthy/standard treatment; treatment group 1 (KP1): 5% sage leaf extract gel application after opening (tissue taken at the 13th minute); and treatment group 2 (KP2): 5% sage leaf extract gel application after opening (tissue taken at the 103rd minute).¹⁶ Data analysis was performed using the Kruskal-Wallis test followed by the Mann-Whitney post hoc test ($p \leq 0.05$), indicating a statistically significant difference.

This research has been conducted in several laboratories of Universitas Jenderal Achmad Yani Cimahi, namely the Animal Experiment

Laboratory of the Faculty of Medicine for the manufacture of punch wound tissue of Wistar rats, the Biochemistry Laboratory of the Faculty of Medicine for the manufacture of sage leaf extract, the Pharmacology Laboratory of the Faculty of Pharmacy for the manufacture of 5% sage leaf extract gel, and the Cytohistotechnology Laboratory of the Faculty of Technology and Health Sciences for the calculation of the number of neutrophils. Ethical clearance has been approved by the Health Research Ethics Committee of the Faculty of Medicine, Universitas Jenderal Achmad Yani, number 019/UH3.09/2023.

This study begins with preparing 5% sage leaf extract gel through the maceration technique. Dried sage leaves are then powdered and sieved to form simplisia, then macerated with 96% ethanol in a ratio (1:9) between simplisia and 96% ethanol leave for 3–5 days. The maceration results are extracted using a rotary evaporator and then dried again in the oven to achieve a consistent consistency. The 5% (w/v) sage leaf extract gel was prepared by dissolving 5 grams of sage leaf powder in solvent to a final volume of 100 ml. The 5% sage leaf extract gel was prepared by dissolving the sage leaf extract in carboxyl methyl cellulose and distilled water, then heating and stirring for 10 minutes to form a 5% sage leaf extract gel. Twenty-five subjects in this study used Wistar strain rats, which were acclimatized for 7 days before treatment. Punch wounds were made under both mandibular anterior teeth with a diameter of 2.5 mm and a depth of 2 mm, previously anesthetized with a CO₂ inhalation concentration of 30–70%.¹⁷ Samples of each group were observed under five fields of view to count the number of neutrophils. The neutrophil counts from each field were then summed to obtain the total count for that sample. Each field of view was divided into several quadrants to facilitate the counting process.

Results

The number of neutrophils in the treatment group is higher than that in the group without any drug or therapy, and the group whose tissue was taken at the 103rd minute has a higher number than the group whose tissue was taken at the 13th minute (Figure).

Table 1 shows that the mean number of




Figure Neutrophil Microscopy

Note: a microscopic picture of 1 quadrant (1,000× magnification) of 1 field of view (400× magnification) in 1 group sample; arrows indicate neutrophils; (A) K-1: without any drug/therapy at 13th minute; (B) K-2: without any drug/therapy at 103rd minute; (C) KN: healthy/normal; (D) KP1: 5% sage leaf extract gel at 13th minute; (E) KP2: 5% sage leaf extract gel at 103rd minute

neutrophils in rats whose tissues were taken at the 13th minute (K-1) without any drug or therapy has a mean of 5.60. In contrast, the 5% sage leaf extract gel (KP1) has a mean of 13.40, which is higher than that without therapy. In rats whose tissues were taken at the 103rd minute (K-2) without receiving any drug or treatment, the mean was 18.40. In contrast, (KP2) sage leaf extract gel 5% had a mean of 22.40, higher than the mean without therapy. In rats (KN) has a mean of 5.40, making it the lowest compared

to other groups. A $p < 0.05$ ($p < 0.000$) indicates a significant difference between groups.

Table 2 shows that group (K-1) has a significant difference with group (K-2), group (KP1), and group (KP2) with $p < 0.05$. Group (K-2) has a significant difference with group (KN) and group (KP1) with $p < 0.05$. There is a significant difference between the group (KN) and the groups (KP1) and (KP2), with $p < 0.05$. Group (KP1) there is a significant difference with a group (KP2) with $p < 0.05$. Group (KP1) showed no significant difference compared to group (KN). Similarly, group (K-2) showed no significant difference compared to group (KP2) at the 103rd minute.

Table 1 Mean Neutrophil Cell Count in the Study Group

Groups	n	Mean	p
K-1	5	5.60	<0.000*
K-2	5	18.40	
KN	5	5.40	
KP1	5	13.40	
KP2	5	22.40	
Total	25	26.08	

Note: Kruskal-Wallis test, *significance $p < 0.05$, K-1: without any drug/therapy at 13th minute, K-2: without any drug/therapy at 103rd minute, KN: healthy/normal, KP1: 5% sage leaf extract gel at 13th minute, KP2: 5% sage leaf extract gel at 103rd minute

Discussion

Table 1 shows that the KP1 and KP2 groups, which were treated with 5% sage leaf extract gel, had a higher average number of neutrophils than the untreated groups (K-1, K-2) and the normal group (KN). This increase is attributed to the bioactive components in the sage leaf extract—namely flavonoids, terpenoids, and rosmarinic acid—which are known to influence neutrophil activity and play significant roles in accelerating

Table 2 Post Hoc Test Analysis between Groups

Groups	K-1	K-2	KN	KP1	KP2
K-1		0.009*	0.916	0.009*	0.009*
K-2	0.009*		0.009*	0.028*	0.076
KN	0.916	0.009*		0.009*	0.009*
KP1	0.009*	0.028*	0.009*		0.009*
KP2	0.009*	0.076	0.009*	0.009*	

Note: Kruskal-Wallis test, *significance $p < 0.05$, K-1: without any drug/therapy at 13th minute, K-2: without any drug/therapy at 103rd minute, KN: healthy/normal, KP1: 5% sage leaf extract gel at 13th minute, KP2: 5% sage leaf extract gel at 103rd minute

the wound healing process. Flavonoids contribute to anti-inflammatory and antioxidant activities, support cell proliferation, and modulate the immune system.^{12–14} These actions influence the proliferation and differentiation of neutrophil cells in the bone marrow and enhance neutrophil recruitment to the wound site.¹⁸ Terpenoids and rosmarinic acid also act as anti-inflammatory, antioxidant, and antimicrobial agents, thereby supporting neutrophil-mediated phagocytosis of bacteria.^{19,20}

The anti-inflammatory properties of these compounds help regulate neutrophil activity by inhibiting the production of pro-inflammatory cytokines and other mediators.^{21,22} However, it is essential to note that excessive neutrophil accumulation may result in tissue damage and chronic inflammation, as neutrophils release proteolytic enzymes that can harm surrounding healthy tissue.²³ In this context, the antioxidant role of rosmarinic acid is particularly valuable, as it inhibits neutrophil adhesion to vascular endothelium and protects tissues from oxidative stress, thus contributing to the controlled regulation of neutrophil levels.^{18,22}

Further analysis of Table 1 indicates that the average number of neutrophils was higher in groups where tissue samples were taken at the 103rd minute compared to those sampled at the 13th minute. Upon injury, the body responds by releasing inflammatory mediators, including neutrophils, which are essential for pathogen clearance through the release of reactive oxygen species (ROS).^{8,9} In humans, neutrophil counts begin to rise around the 6th hour post-injury and peak within 48 hours. When converted to an equivalent timeline in mice, this corresponds to the period between the 13th and 103rd minutes post-injury. This increase occurs due to

pathogenic bacteria that have toxic properties; they can stimulate an inflammatory response.^{10,24} The higher neutrophil count observed at the 103rd minute reflects this physiological accumulation, combined with chemotaxis—the chemical signaling that directs neutrophils rapidly to the wound site. These findings align with Dutta and Sengupta's¹⁰ rat-to-human age conversion model, which supports a 28:1 ratio for physiological timelines.

Table 2 shows that the mean neutrophil counts between the K-1 and KN groups were not significantly different ($p = 0.916$). The presence of neutrophils in non-wounded tissues, such as in the KN group, is because neutrophils are the most abundant circulating leukocytes in the bloodstream, making them detectable even without injury. Additionally, factors such as physiological stress, allergic responses, or latent infections may promote neutrophil migration in the absence of visible wounds.²⁴ The similarity in neutrophil count between K-1 and KN is also consistent with the fact that the 13th minute represents the very early phase of the inflammatory response, during which tissue damage is still being detected, and vasodilation is just beginning to occur.^{10,25}

Meanwhile, the comparison between the K-2 and KP2 groups also showed no statistically significant difference in neutrophil count ($p = 0.076$). In the K-2 group, the absence of therapeutic intervention with sage leaf extract meant that neutrophil activity remained relatively unregulated. Without the anti-inflammatory and antioxidant effects of flavonoids, terpenoids, and rosmarinic acid, neutrophils in the K-2 group exhibited slower phagocytic function, oxidative stress in surrounding tissues, and delayed apoptosis. These factors may contribute to an

accumulation of neutrophils in the wound area, similar to what was observed in the KP2 group, although for different reasons.^{20,22}

Overall, the findings suggest that 5% sage leaf extract gel positively influences neutrophil activity in the wound-healing process of Wistar rats. It is consistent with previous research, which demonstrated that sage leaf extract—particularly at a concentration of 5%—can significantly accelerate wound healing and holds promise for clinical applications in wound care.¹⁵ The increase in neutrophil count appears to play a critical role in this process, enhancing bacterial clearance through phagocytosis while remaining within a regulated range to avoid tissue damage. Thus, the bioactive compounds in sage leaf extract not only promote neutrophil recruitment but also help maintain a balanced inflammatory response during healing.²⁶

Conclusions

This study concluded that 5% sage leaf extract gel (*Salvia officinalis* L.) has the potential to increase neutrophil count during the wound healing process in Wistar rats with punch-induced injuries, thereby supporting and accelerating the healing process. With further research and development, this extract is a promising alternative compound for clinical use in wound care.

Conflict of Interest

The authors declare that there is no conflict of interest.

Acknowledgment

We want to thank everyone who helped and supported this research, starting from the Biochemistry Laboratory of the Faculty of Medicine, Animal Experiment Laboratory of the Faculty of Medicine, Pharmacology Laboratory of the Faculty of Pharmacy, Cytohistotechnology Laboratory of the Faculty of Technology and Health Sciences, Universitas Jenderal Achmad Yani, Cimahi, and also IP Farms Lembang.

References

1. Firdaus NZ, Alda AA, Gunawan IS. Potensi kandungan biji anggur dalam mempercepat penyembuhan luka. JPPP. 2020;2(2):139–46.
2. Wintoko R, Yadika ADN. Manajemen terkini perawatan luka. JK Unila. 2020;4(2):183–9.
3. Montazer Lotf-Elahi MS, Farzinnia G, Jaafari-Ashkavandi Z. Clinicopathological study of 1000 biopsied gingival lesions among dental outpatients: a 22-year retrospective study. BMC Oral Health. 2022;22(1):154.
4. Sitorus EP, Julianto I. Teknik-teknik biopsi kulit. CDK. 2018;45(6):466–71.
5. Serra MB, Barroso WA, da Silva NN, Silva SDN, Borges ACR, Abreu IC, et al. From inflammation to current and alternative therapies involved in wound healing. Int J Inflam. 2017;2017:3406215.
6. Zhao H, Wu L, Yan G, Chen Y, Zhou M, Wu Y, et al. Inflammation and tumor progression: signaling pathways and targeted intervention. Signal Transduct Target Ther. 2021;6(1):263.
7. Gierlikowska B, Stachura A, Gierlikowski W, Demkow U. Phagocytosis, degranulation and extracellular traps release by neutrophils—the current knowledge, pharmacological modulation and future prospects. Front Pharmacol. 2021;12:666732.
8. Cui LL, Zhang Y, Chen ZY, Su YY, Liu Y, Boltze J. Early neutrophil count relates to infarct size and fatal outcome after large hemispheric infarction. CNS Neurosci Ther. 2020;26(8):829–36.
9. Bo L, Guojun T, Li G. An expanded neuroimmunomodulation axis: sCD83-indoleamine 2,3-dioxygenase-kynurenine pathway and updates of kynurenine pathway in neurologic diseases. Front Immunol. 2018;9:1363.
10. Dutta S, Sengupta P. Men and mice: relating their ages. Life Sci. 2016;152:244–8.
11. Purwanto E, Rajab MA, Angraini R. Potensi kandungan bahan-bahan alami untuk proses penyembuhan luka: literatur review. J Ilm Keperawatan Kebidanan Holistic Care. 2023;6(2):18–25.
12. Slipranata M, Lestari FB, Sandi NA, Salasia SIO. Potensi ekstrak daun sage (*Salvia officinalis* L.) sebagai anti-*Streptococcus suis* penyebab zoonotik meningitis. J Sain Vet. 2016;34(2):198–202.
13. Blezeinsky FN, Gumay AR, Hardian. Efek pemberian ekstrak daun *Carica pubescens*

- terhadap jumlah neutrofil pada tikus Sprague Dawley yang diinduksi azoxymethane. JKD. 2019;8(3):955–63.
14. Hikmah U, Triastuti A. Mechanism and immunomodulator bioactive compounds of *Phyllanthus niruri* (meniran). JIF. 2022;18(2):205–18.
 15. Karimzadeh S, Farahpour MR. Topical application of *Salvia officinalis* hydroethanolic leaf extract improves wound healing process. Indian J Exp Biol. 2017;55(2):98–106.
 16. Dahlan MS. Metode MSD: pintu gerbang memahami epidemiologi, biostatistik, dan metode penelitian. 2nd Edition. Jakarta: Sagung Seto; 2017.
 17. Chisholm JM, Pang DS. Assessment of carbon dioxide, carbon dioxide/oxygen, isoflurane and pentobarbital killing methods in adult female Sprague-Dawley rats. PLoS One. 2016;11(9):e0162639.
 18. Dahliani R, Nahzi MYI, Setyawardhana RHD. Pengaruh ekstrak daun karamunting (*Rhodomyrtus tomentosa*) terhadap jumlah sel neutrofil pada pulpa. Dentin. 2021; 5(3):122–8.
 19. Tousif MI, Nazir M, Riaz N, Saleem M, Tauseef S, Azam SM, et al. Terpenoids as human neutrophil elastase (HNE) inhibitors: a comprehensive review of natural anti-inflammatory isoprenoids. Chembiochem. 2023;24(22):e202300346.
 20. Azalia D, Rachmawati I, Zahira S, Andriyani F, Sanini TM, Supriyatin, et al. Uji kualitatif senyawa aktif flavonoid dan terpenoid pada beberapa jenis tumbuhan Fabaceae dan Apocynaceae di kawasan TNGPP Bodogol. Bioma. 2023;8(1):32–43.
 21. Hamidah M, Moektiwardoyo M, Abdassah M. Review artikel: senyawa aktif antiinflamasi daun jawer kotok (*Plectranthus scutellarioides* (L.) R.Br.). Farmaka. 2019;17(1):89–96.
 22. Prima W, Samsu N, Khotimah H. Comparison of the effect of vitamin E and rosmarinic acid on expression of p65 NF- κ B subunit in diabetes rat. AMS. 2020;6(2):117–23.
 23. Li Y, Wang W, Yang F, Xu Y, Feng C, Zhao Y. The regulatory roles of neutrophils in adaptive immunity. Cell Commun Signal. 2019;17(1):147.
 24. Pribadi N, Rahayu RP, Ismiyatin K, Putri CR, Surboyo MDC. The effect of lipoteichoic acid from *Lactobacillus plantarum* on dental pulp inflammation. Eur J Dent. 2021;15(4):682–6.
 25. Faisal SF, Ingriyani CG, Mulia VD. Hubungan jumlah neutrofil dengan umur luka memar pada mencit (*Mus musculus*) strain Balb-C. JIM Kedokteran Biomedis. 2018;3(1):8–13.
 26. Raziyeve K, Kim Y, Zharkinbekov Z, Kassymbek K, Jimi S, Saparov A. Immunology of acute and chronic wound healing. Biomolecules. 2021;11(5):700.